




Turf Products

0.72G Prophecy®

On DG^{PRO}



Dispersing Granule Technology

- Broad spectrum and systemic disease control and suppression for turf
- Dispersing granule fungicide
- Easy to spread formula

Active Ingredient: Propiconazole 0.72%

Other ingredients: 99.28%

Total 100.00%

*CAS No. 60207-90-1
EPA Reg. No. 9198-219
EPA EST. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A, 9198-IN-3S
Underlined letter is first letter used in run code on bag

KEEP OUT OF REACH OF CHILDREN
CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements
Hazards to Humans and Domestic Animals
CAUTION
Causes moderate eye irritation. Harmful if swallowed, absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear goggles or face shield. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment
Applicators and other handlers must wear:
• Long-sleeved shirt and long pants
• Chemical resistant gloves made from any waterproof material
• Shoes plus socks
• Goggles or face shield

USER SAFETY REQUIREMENTS
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations
Users should:
• Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards
This pesticide is toxic to fish and shrimp. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment, disposing of equipment wash waters or rinsate.

FIRST AID
If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
Note to Physician: If ingested, induce emesis or lavage stomach. Treat symptomatically.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-9851 for emergency medical treatment information.

Net Wt. 25 Lbs. (11.34 kg)



The Andersons Turf Products 0.72G Prophecy on DG PRO

The Andersons Turf Products 0.72G Prophecy on DG^{PRO}

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons or pets, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Workers Protection Standard 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: Coveralls, shoes and socks, and chemical-resistant gloves made from any waterproof material.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to use of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The Worker Protection Standard applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow adults, children, or pets to enter the treated areas until dust has settled.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store in a clean, dry secure place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Pesticide wastes may be acutely hazardous. Improper disposal is a violation of Federal law. Pesticides, mixtures, or equipment rinse water that cannot be chemically reprocessed must be disposed of according to applicable Federal, state or local procedures. Contact your state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

General Information

This product is a systemic fungicide for use on turfgasses for the control and suppression of:

Dollar spot	Sclerotinia homoeocarpa	Spring dead spot	Leptophyllum kossae, Leptophyllum nammat, Ophiophthalmia heptochloa, Gaumannomyces graminis
Brown patch	Rhizoctonia solani	Take-all patch	Gaumannomyces graminis
Anthraxnose	Colletotrichum graminicola	Leafspot	Biopolaris spp., Drechslera spp.
Red thread	Lasiolesia fuctiformis	Gray Leafspot	Pyricularia grisea
Pink patch	Lasiolesia roseopellis	Pink snowmold	Microdochium nivale
Rust	Puccinia graminis	Fusarium patch	Microdochium nivale
Powdery mildew	Erysiphe graminis	Gray Snowmold	Typhula spp.
Stripe smut	Ustilago striiformis and Ustilago agrostii	Yellow patch	Rhizoctonia cerealis
Summer patch	Magnaporthe poae	Zyzybia patch	Rhizoctonia solani
Neotrophic ring spot	Leptophyllum kossae		

Do not apply this product through any type of irrigation system.

Use this product in a preventative disease control program. Properly calibrate the spreader to ensure thorough coverage. Apply after mowing. For control of foliar diseases, apply to wettest foliage. For control of soil-borne diseases, this product can be watered in after application. Under conditions optimum for high disease pressure, use the higher rate and the shorter interval. For optimum turf quality and disease control, use this product in conjunction with turf management practices that promote good plant health and optimum disease control. Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.

Do not apply more than 20 lbs. product/1,000 sq. ft./calendar year.

Important: Bermudagrass can be sensitive to this product. Do not exceed 5 lbs./1,000 sq. ft. every 30 days on any variety of bermudagrass. In Florida, do not apply this product to bermudagrass golf greens when temperatures exceed 90° F.

Note: Do not graze animals on treated areas. Do not feed clippings from treated areas to livestock or poultry.



Turfgrass - Specific Diseases, Rates, and Application Timing

Disease	Lbs. Product Per 1,000 Sq. Ft.	Lbs. Product Per Acre	Application Timing/Interval	Instructions
Dollar spot <i>Sclerotinia homoeocarpa</i>	2.5	100	14-28 days	Make no more than 3 consecutive applications for dollar spot before rotating to an alternative EPA-registered fungicide having a different mode of action.
Anthraxnose <i>Colletotrichum graminicola</i>	2.5	100	14-28 days	Apply when conditions are favorable for disease development. When disease pressure is high, use shorter intervals.
Brown Patch <i>Rhizoctonia solani</i>	2.5	100	14-21 days	Begin applications in May or June before disease is present. Under conditions of high temperatures and high humidity, use this product at shorter intervals.
Powdery mildew <i>Erysiphe graminis</i> Rust <i>Puccinia graminis</i>	2.5	100	14-28 days	Apply when conditions are favorable for disease development.
Red thread <i>Lasiolesia fuctiformis</i> Pink patch <i>Lasiolesia roseopellis</i>	2.5	100	14-21 days	Apply when conditions are favorable for disease development.
Stripe smut <i>Ustilago striiformis</i> <i>Ustilago agrostii</i>	2.5	100	Fall or spring	Apply once in the fall after grass becomes dormant or in the early spring before grass starts to grow.
Gray leafspot <i>Pyricularia grisea</i>	2.5	100	14 days	Apply when conditions are favorable for disease development.
Melting Out Leafspot <i>Empoisma spp.</i> <i>Drechslera spp.</i>	2.5	100	14 days	Under light to moderate pressure, apply this product to reduce the severity of leaf spot and melting out caused by Helminthosporium-type pathogens.
Summer patch <i>Magnaporthe poae</i>	2.5-5	100-216	14 days 20 days	Apply this product beginning in April. Use the 5 lbs./1,000 sq. ft. rate on a 20 day schedule and the 2.5 lbs./1,000 sq. ft. rate on a 14 day schedule.
Take-all patch <i>Gaumannomyces graminis</i>	2.5-5	100-216	Spring and Fall	Apply this product to reduce the severity of take-all patch. Make 1-2 fall applications in September and October or when night temperatures drop to 55° F and 1-2 spring applications in April and May, depending on local recommendations.
Spring dead spot <i>Leptophyllum kossae</i> <i>Leptophyllum nammat</i> <i>Ophiophthalmia heptochloa</i> <i>Gaumannomyces graminis</i>	5	216	30 days	Make 1-3 applications. For one application, apply in September or October. For multiple applications, begin applications in August.
Neotrophic ring spot <i>Leptophyllum kossae</i>	5	216	Fall or spring	Apply in the fall and/or the early spring depending on local recommendations.
Snowmold Gray <i>Typhula spp.</i> Pink <i>Microdochium nivale</i>	2.5-5	100-216	Late Fall	Apply one application in the late fall before snow cover. Do not apply on top of snow. For optimum disease control, the 2.5 and 3.75 lbs. rates should be used in conjunction with either PCOB or chlorothalonil at label rates.
Fusarium patch <i>Microdochium nivale</i>	2.5-5	100-216	Fall-Early Spring	Apply when conditions are favorable for disease development.
Yellow patch <i>Rhizoctonia cerealis</i>	3.75-5	163-216	Late Fall	Apply one application in the late fall before snow cover. Do not apply on top of snow. If using a 3.75 lb./1,000 sq. ft. rate, use in conjunction with a registered contact fungicide at the label rate.
Zyzybia patch, Large patch of zyzybia <i>Rhizoctonia solani</i>	3.75-5	163-216	Early Fall	Make one application in the early fall (mid-September to mid-October) prior to development of disease symptoms. Consult local turfgrass extension agents to determine the optimum application timing in your area.

Dichondra - Specific Diseases, Rates, and Application Timing

Disease	Lbs. Product Per 1,000 Sq. Ft.	Lbs. Product Per Acre	Application Timing/Interval	Instructions
Dichondra Rust <i>Puccinia dichondrae</i>	2.5	100	14-21 days	Apply when conditions are favorable for disease development.

Suggested Spreader Settings

These suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product.
25 lbs. treats 10,000 sq. ft. (0.23 acre) at the 100 lbs/acre rate (2.20 lbs./1,000 sq. ft. rate)
25 lbs. treats 6,000 sq. ft. (0.15 acre) at the 163 lbs/acre rate (3.75 lbs./1,000 sq. ft. rate)
25 lbs. treats 5,000 sq. ft. (0.11 acre) at the 216 lbs/acre rate (5.00 lbs./1,000 sq. ft. rate)

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	RECOMMENDED RATE per 1,000 sq. ft.		
			2.5W (0.78 lbs. ai/acre)	3.75W (1.17 lbs. ai/acre)	5.00W (1.57 lbs. ai/acre)
Andersons Model 2000 (zone 4); Andersons 2000 SR (zone 4); Andersons LCO-1000	3 mph	9 feet	J	K 1/4	L 1/2
Andersons SSD /SS-2 Drop	3 mph	Overlap wheels	4 1/2	5 1/2	6 1/2
Scotts R-8, R-8A (zone 4)	3 mph	9 feet	H 1/2	I 3/4	K
LESCO 20000 (system 0.25)	3 mph	9 feet	D	D 1/2	E
LESCO 70000 (system 0.25)	3 mph	6 feet	12	11	12
Law Mower WPR, WPR, HR, 1250 (PTO at 450 rpm)	4.5 mph	30 feet	4 B/A	4 3/4 B/A	5 1/2 B/A
Vicon (03 Series)	4.5 mph	21 feet	13	16	20

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION OF LIABILITY: Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS: THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS. YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE JURISDICTION TO STATE JURISDICTION.

Manufactured by:
The Andersons, Inc.
P.O. Box 119
Mumme, OH 43537

Visit our website: www.andersonsplantnutrient.com

©2019 The Andersons, Inc.
Prophecy is a trademark of The Andersons Inc.; DG Pro and The Andersons logo are registered trademarks of The Andersons, Inc. 091919

APTDPG22
APTDPG22-1B

